

13-06-2001  
X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)

P DESCAMD

-1-

## Novel sPLA<sub>2</sub> Inhibitors

### Field of the Invention

This invention relates to novel indole compounds  
5 useful for Inflammatory Diseases.

### Background of the Invention

The structure and physical properties of human non-  
pancreatic secretory phospholipase A<sub>2</sub> (hereinafter  
10 called, "sPLA<sub>2</sub>") has been thoroughly described in two  
articles, namely, "Cloning and Recombinant Expression of  
Phospholipase A<sub>2</sub> Present in Rheumatoid Arthritic  
Synovial Fluid" by Seilhamer, Jeffrey J.; Pruzanski,  
Waldemar; Vadas Peter; Plant, Shelley; Miller, Judy A.;  
15 Kloss, Jean; and Johnson, Lorin K.; The Journal of  
Biological Chemistry, Vol. 264, No. 10, Issue of April  
5, pp. 5335-5338, 1989; and "Structure and Properties of  
a Human Non-pancreatic Phospholipase A<sub>2</sub>" by Kramer, Ruth  
M.; Hession, Catherine; Johansen, Berit; Hayes,  
20 Gretchen; McGray, Paula; Chow, E. Pingchang; Tizard,  
Richard; and Pepinsky, R. Blake; The Journal of  
Biological Chemistry, Vol. 264, No. 10, Issue of April  
5, pp. 5768-5775, 1989; the disclosures of which are  
incorporated herein by reference.

25

-2-

It is believed that sPLA<sub>2</sub> is a rate limiting enzyme in the arachidonic acid cascade which hydrolyzes membrane phospholipids. Thus, it is important to develop compounds which inhibit sPLA<sub>2</sub> mediated release of fatty acids (e.g., arachidonic acid). Such compounds would be of value in general treatment of conditions induced and/or maintained by overproduction of sPLA<sub>2</sub>; such as sepsis or rheumatoid arthritis.

It is desirable to develop new compounds and treatments for sPLA<sub>2</sub> induced diseases.

#### Summary of the Invention

This invention provides novel indole compounds having potent and selective effectiveness as inhibitors of mammalian sPLA<sub>2</sub>.

This invention is also the use of novel indole compounds useful in the treatment and prevention of Inflammatory Diseases.

This invention is also the use of novel of indole compounds to inhibit mammalian sPLA<sub>2</sub> mediated release of fatty acids.

25

-3-

This invention is also a pharmaceutical composition containing any of the indole compounds of the invention.


5    **I.    Definitions:**

          The term, "Inflammatory Diseases" refers to  
diseases such as inflammatory bowel disease, sepsis,  
septic shock, adult respiratory distress syndrome,  
pancreatitis, trauma-induced shock, bronchial asthma,  
10    allergic rhinitis, rheumatoid arthritis, cystic  
fibrosis, stroke, acute bronchitis, chronic bronchitis,  
acute bronchiolitis, chronic bronchiolitis,  
osteoarthritis, gout, spondylarthropathris, ankylosing  
spondylitis, Reiter's syndrome, psoriatic arthropathy,  
15    enteropathic spondylitis, Juvenile arthropathy or  
juvenile ankylosing spondylitis, Reactive arthropathy,  
infectious or post-infectious arthritis, gonococcal  
arthritis, tuberculous arthritis, viral arthritis,  
fungal arthritis, syphilitic arthritis, Lyme disease,  
20    arthritis associated with "vasculitic syndromes",  
polyarteritis nodosa, hypersensitivity vasculitis,  
Luegenec's granulomatosis, polymyalgin rheumatica, joint  
cell arteritis, calcium crystal deposition  
arthropathris, pseudo gout, non-articular rheumatism,  
25    bursitis, tenosynovitis, epicondylitis (tennis elbow),  
carpal tunnel syndrome, repetitive use injury (typing).

-4-

miscellaneous forms of arthritis, neuropathic joint  
disease (charco and joint), hemarthrosis (hemarthrosic),  
Henoch-Schonlein Purpura, hypertrophic osteoarthropathy,  
multicentric reticulohistiocytosis, arthritis associated  
5 with certain diseases, surcoilosis, hemochromatosis,  
sickle cell disease and other hemoglobinopathries,  
hyperlipoproteineimia, hypogammaglobulinemia,  
hyperparathyroidism, acromegaly, familial Mediterranean  
fever, Behat's Disease, systemic lupus erythrematosis,  
10 or relapsing polychondritis and related diseases which  
comprises administering to a mammal in need of such  
treatment a therapeutically effective amount of the  
compound of formula I in an amount sufficient to inhibit  
sPLA<sub>2</sub> mediated release of fatty acid and to thereby  
15 inhibit or prevent the arachidonic acid cascade and its  
deleterious products.

The term, "indole nucleus" refers to a nucleus  
(having numbered positions) with the structural  
20 formula (X):



The indole compounds of the invention employ  
certain defining terms as follows:

-5-

The term, "alkyl" by itself or as part of another  
substituent means, unless otherwise defined, a straight  
or branched chain monovalent hydrocarbon radical such as  
methyl, ethyl, n-propyl, isopropyl, n-butyl, tertiary  
5 butyl, sec-butyl, n-pentyl, and n-hexyl.

The term, "alkenyl" employed alone or in  
combination with other terms means a straight chain or  
branched monovalent hydrocarbon group having the stated  
10 number range of carbon atoms, and typified by groups  
such as vinyl, propenyl, crotonyl, isopentenyl, and  
various butenyl isomers.

The term, "hydrocarbyl" means an organic group  
15 containing only carbon and hydrogen.

The term, "halo" means fluoro, chloro, bromo, or  
iodo. The term, heterocyclic radical, refers to radicals  
derived from monocyclic or polycyclic, saturated or  
20 unsaturated, substituted or unsubstituted heterocyclic  
nuclei having 5 to 14 ring atoms and containing from 1 to  
3 hetero atoms selected from the group consisting of  
nitrogen, oxygen or sulfur. Typical heterocyclic radicals  
are pyrrolyl, pyrrolodiny, piperidiny, furanyl,  
25 thiophenyl, pyrazolyl, imidazolyl, phenylimidazolyl,  
triazolyl, isoxazolyl, oxazolyl, thiazolyl, thiadiazolyl,


-6-

indolyl, carbazolyl, norharmanyl, azaindolyl,  
benzofuranyl, dibenzofuranyl, dibenzothiophenyl,  
indazolyl, imidazo(1.2-A)pyridinyl, benzotriazolyl,  
anthranilyl, 1,2-benzisoxazolyl, benzoxazolyl,  
5 benzothiazolyl, purinyl, pyridinyl, dipyridyl,  
phenylpyridinyl, benzylpyridinyl, pyrimidinyl,  
phenylpyrimidinyl, pyrazinyl, 1,3,5-triazinyl, quinolinyl,  
phthalazinyl, quinazolinyl, morpholino, thiomorpholino,  
homopiperazinyl, tetrahydrofuranyl, tetrahydropyranyl,  
10 oxacanyl, 1,3-dioxolanyl, 1,3-dioxanyl, 1,4-dioxanyl,  
tetrahydrothiophenyl, pentamethylenesulfadyl, 1,3-  
dithianyl, 1,4-dithianyl, 1,4-thioxanyl, azetidyl,  
hexamethyleneiminium, heptamethyleneiminium, piperazinyl  
and quinoxalinyl.

15

The term, "carbocyclic radical" refers to radicals  
derived from a saturated or unsaturated, substituted or  
unsubstituted 5 to 14 membered organic nucleus whose ring  
forming atoms (other than hydrogen) are solely carbon  
20 atoms. Typical carbocyclic radicals are cycloalkyl,  
cycloalkenyl, phenyl, spiro[5.5]undecanyl, naphthyl,  
norbornanyl, bicycloheptadienyl, tolulyl, xylenyl,  
indenyl, stilbenyl, terphenyl, diphenylethylenyl,  
phenyl-cyclohexenyl, acenaphthylenyl, and anthracenyl,  
25 biphenyl, bibenzyl and related bibenzyl homologues  
represented by the formula (a):

-7-



where n is a number from 1 to 8.

5        The term, "non-interfering substituent", refers to radicals suitable for substitution at positions 4,5,6 and/or 7 of the indole nucleus and on other nucleus substituents (as hereinafter described for Formula I), and radicals suitable for substitution on the

10   heterocyclic radical and carbocyclic radical as defined above. Illustrative non-interfering radicals are C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>2</sub>-C<sub>8</sub> alkenyl, C<sub>2</sub>-C<sub>8</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> aralkyl, C<sub>7</sub>-C<sub>12</sub> alkaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, phenyl, tolulyl, xylenyl, biphenyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>2</sub>-C<sub>8</sub>

15   alkenyloxy, C<sub>2</sub>-C<sub>8</sub> alkynyloxy, C<sub>2</sub>-C<sub>12</sub> alkoxyalkyl, C<sub>2</sub>-C<sub>12</sub> alkoxyalkyloxy, C<sub>2</sub>-C<sub>12</sub> alkylcarbonyl, C<sub>2</sub>-C<sub>12</sub> alkylcarbonylamino, C<sub>2</sub>-C<sub>12</sub> alkoxyamino, C<sub>2</sub>-C<sub>12</sub> alkoxyaminocarbonyl, C<sub>1</sub>-C<sub>12</sub> alkylamino, C<sub>1</sub>-C<sub>6</sub> alkylthio, C<sub>2</sub>-C<sub>12</sub> alkylthiocarbonyl, C<sub>1</sub>-C<sub>8</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>8</sub>


20   alkylsulfonyl, C<sub>2</sub>-C<sub>8</sub> haloalkoxy, C<sub>1</sub>-C<sub>8</sub> haloalkylsulfonyl, C<sub>2</sub>-C<sub>8</sub> haloalkyl, C<sub>1</sub>-C<sub>8</sub> hydroxyalkyl, -C(O)O(C<sub>1</sub>-C<sub>8</sub> alkyl), -(CH<sub>2</sub>)<sub>n</sub>-O-(C<sub>1</sub>-C<sub>8</sub> alkyl), benzyloxy, phenoxy, phenylthio, -(CONHSO<sub>2</sub>R), -CHO, amino, amidino,

-8-

bromo, carbamyl, carboxyl, carbalkoxy,  $-(CH_2)_n-CO_2H$ ,  
chloro, cyano, cyanoguanidiny, fluoro, guanidino,  
hydrazide, hydrazino, hydrazido, hydroxy, hydroxyamino,  
iodo, nitro, phosphono,  $-SO_3H$ , thioacetal, thiocarbonyl,  
5 and carbonyl; where n is from 1 to 8 and R is  $C_1-C_8$   
alkyl.

The term, "organic substituent" refers to a  
monovalent radical consisting of carbon and hydrogen  
10 with or without oxygen, nitrogen, sulfur, halogen, or  
other elements. Illustrative organic substituents are  
 $C_1-C_8$  alkyl, aryl,  $C_7-C_{14}$  aralkyl,  $C_7-C_{14}$  alkaryl,  $C_3-C_8$   
cycloalkyl,  $C_1-C_8$  alkoxyalkyl and these groups  
substitued with halogen,  $-CF_3$ ,  $-OH$ ,  $C_1-C_8$  alkyl, amino,  
15 carbonyl, and  $-CN$ .

The term, "acylamino acid group" is represented by  
the formula:



wherein  $R^{4a}$  is selected from the group consisting of H,  
 $(C_1-C_6)$ alkyl,  $(C_1-C_6)$ alkoxy, heteroaryl and aryl,  $-CF_3$ ;  
and wherein  $NR^{4b}$  is an amino acid residue with the



-9-


nitrogen atom being part of the amino group of the amino  
acid. A typical amino acid is selected from the group  
comprising isoleucine, valine, phenylalanine, aspartic  
acid, leucine, glycine, asparagine, cystein, glutamine,  
5 glutamic acid, histidine, lysine, methionine, serine,  
threonine, tryptophan, tyrosine and derivatives thereof.  
Also contemplated within the definition of amino acid is  
l-proline, d-proline and derivatives thereof. Also  
contemplated within the definition of amino acids are  
10 peptides, polypeptides and derivatives thereof.

The term "substituted group" is an organic group  
substituted with one or more non-interfering  
substituents.

15


The terms, "amino acid residue" refer to the  
portion of the amino acid group coupled at the nitrogen  
atom of the amino terminus. It is the amino acid less a  
hydrogen atom from the amino terminus. It is further  
20 illustrated as used herein for the amino acid alanine  
attached at the nitrogen atom as shown below:

-10-






The words, "acylamino acid linker" refer to a divalent linking group symbolized as,  $-(L_C)-$ , which has the function of joining the 4 - position of the indole nucleus to an acylamino acid group in the general

5 relationship:



The words, "acylamino acid linker length", refer to the number of atoms (excluding hydrogen) in the shortest chain of the linking group  $-(L_C)-$  that connects the 4 - position of the indole nucleus with the acylamino acid group. The presence of a carbocyclic ring in  $-(L_C)-$  counts as the number of atoms approximately equivalent to the calculated diameter of the carbocyclic ring. Thus, a benzene or cyclohexane ring in the acid linker counts as 2 atoms in calculating the length of  $-(L_C)-$ . Illustrative acylamino acid linker groups are;

-11-



wherein, groups (a), (b) and (c). have acid linker lengths of 5, 7, and 2, respectively.

5

The term, "(acidic group)" means an organic group which when attached to an indole nucleus at position 5, through suitable linking atoms (hereinafter defined as the "acid linker"), acts as a proton donor capable of hydrogen bonding. Illustrative of an (acidic group) are the following:

-5-tetrazolyl,

-SO<sub>3</sub>H,




13-06-2001

X-12420



00944673.3(11-07-2000) - US00/16319(11-07-2000)

P DESCAMD


-12-



5




-13-



where n is 1 to 8, R<sub>80</sub> is a metal or C<sub>1</sub>-C<sub>8</sub> and R<sub>81</sub>  
 5 is an organic substituent or -CF<sub>3</sub>.




The words, "acid linker" refer to a divalent  
 linking group symbolized as, -(L<sub>a</sub>)-, which has the  
 function of joining the 5 position of the indole nucleus  
 10 to an acidic group in the general relationship:



The words, "acid linker length", refer to the number  
 of atoms (excluding hydrogen) in the shortest chain of the  
 15 linking group -(L<sub>a</sub>)- that connects the 5 position of the

-14-

indole nucleus with the acidic group. The presence of a carbocyclic ring in  $-(L_a)-$  counts as the number of atoms approximately equivalent to the calculated diameter of the carbocyclic ring. Thus, a benzene or cyclohexane ring in  
 5 the acid linker counts as 2 atoms in calculating the length of  $-(L_a)-$ . Illustrative acid linker groups are;



10 wherein, groups (a), (b), and (c) have acid linker lengths of 5, 7, and 2, respectively.

The term, "amine", includes primary, secondary and tertiary amines.

-15-

The terms, "mammal" and "mammalian" include human and domesticated quadrupeds.

The term, "alkylene chain of 1 or 2 carbon atoms" refers to the divalent radicals,  $-\text{CH}_2-\text{CH}_2-$  and  $-\text{CH}_2-$ .

5

The term, "group containing 1 to 4 non-hydrogen atoms" refers to relatively small groups which form substituents at the 2 position of the indole nucleus, said groups may contain non-hydrogen atoms alone, or non-hydrogen atoms plus hydrogen atoms as required to satisfy the unsubstituted valence of the non-hydrogen atoms, for example; (i) groups absent hydrogen which contain no more than 4 non-hydrogen atoms such as  $-\text{CF}_3$ ,  $-\text{Cl}$ ,  $-\text{Br}$ ,  $-\text{NO}_2$ ,  $-\text{CN}$ ,  $-\text{SO}_3$ ; and (ii) groups having hydrogen atoms which contain less than 4 non-hydrogen atoms such as  $-\text{CH}_3$ ,  $-\text{C}_2\text{H}_5$ , and  $-\text{CH}=\text{CH}_2$ .


20

The term "oxime amide" means the radical,  
 $-\text{C}=\text{NOR}-\text{C}(\text{O})\text{NH}_2$

The term "thio-oxime amide" means the radical  
 $-\text{C}=\text{NOR}-\text{C}(\text{S})-\text{NH}_2$ .


The term "spiro[5.5]undecanyl" refers to the group  
represented by the formula;

-16-



## II. The amino acid 1H-indole Compounds of the Invention:

The present invention provides novel classes of  
5 indole compounds useful as sPLA2 inhibitors for the  
treatment of inflammation. Classes of indole compounds  
of this invention include indole glyoxylamide amino acid  
derivatives, indole-3-oxime amide amino acid derivatives  
and indole acetamide amino acid derivatives. The  
10 compounds of the invention have the general formula (I)  
or a pharmaceutically acceptable salt, solvate or  
prodrug thereof;



wherein ;

R<sub>1</sub> is selected from groups (a), (b), and (c)

wherein;



-17-


(a) is C7-C20 alkyl, C7-C20 haloalkyl, C7-C20 alkenyl, C7-C20 alkynyl, carbocyclic radical, or heterocyclic radical, or

(b) is a member of (a) substituted with one or  
5 more independently selected non-interfering substituents; or

(c) is the group  $-(L_1)-R_{11}$ ; where,  $-(L_1)-$  is a divalent linking group of 1 to 8 atoms and where  $R_{11}$  is a group selected from (a) or (b);

10  $R_2$  is hydrogen, or a group containing 1 to 4 non-hydrogen atoms plus any required hydrogen atoms;


$R_3$  is  $-(L_3)-Z$ , where  $-(L_3)-$  is a divalent linker group selected from a bond or a divalent group selected from:




15

and Z is selected from an oxime amide or oxime thioamide group represented by the formulae,


-18-



or



or



5

wherein  $\text{X}$  is oxygen or sulfur,  $\text{R}_a$  is independently selected from hydrogen,  $\text{C}_1$ - $\text{C}_8$  alkyl, aryl,  $\text{C}_1$ - $\text{C}_8$  alkaryl,  $\text{C}_1$ - $\text{C}_8$  alkoxy, aralkyl and  $-\text{CN}$ ;

$\text{R}_4$  is the group,  $-(\text{L}_c)-(\text{acylamino acid group})$ ;

10 wherein  $-(\text{L}_c)-$ , is an acylamino acid linker having an acylamino acid linker length of 1 to 8;

$\text{R}_5$  is selected from hydrogen, a non-interfering substituent, or the group,  $-(\text{L}_a)-(\text{acidic group})$ ; wherein  $-(\text{L}_a)-$ , is an acid linker having an acid linker length

15 of 1 to 8.

-19-

R<sub>6</sub> and R<sub>7</sub> are selected from hydrogen, non-interfering substituent, carbocyclic radical, carbocyclic radical substituted with non-interfering substituent(s), heterocyclic radicals, and heterocyclic radical substituted with non-interfering substituent(s).

**Preferred Subgroups of Compounds of Formula (I):**  
**Preferred R<sub>1</sub> substituents:**


A preferred subclass of compounds of formula (I) are those where for R<sub>1</sub> the divalent linking group -(L<sub>1</sub>)- is a group represented by any one of the following formulae (Ia), (Ib), (Ic), (Id), (Ie), or (If):



15

or

-20-



where  $Q_1$  is a bond or any of the divalent groups (Ia),  
(Ib), (Ic), (Id), (Ie), and (If) and each  $R_{10}$  is


5 independently hydrogen,  $C_{1-8}$  alkyl,  $C_{1-8}$  haloalkyl or  
 $C_{1-8}$  alkoxy.

Particularly preferred as the linking group  $-(L_1)-$  of  
 $R_1$  is an alkylene chain of 1 or 2 carbon atoms, namely,  
10  $-(CH_2)-$  or  $-(CH_2-CH_2)-$ .

The preferred group for  $R_{11}$  is a substituted or  
unsubstituted group selected from the group consisting of  
 $C_5-C_{14}$  cycloalkyl,  $C_5-C_{14}$  cycloalkenyl, phenyl, naphthyl,  
15 norbornanyl, bicycloheptadienyl, tolulyl, xylenyl,  
indenyl, stilbenyl, terphenyl, diphenylethylenyl,  
phenyl-cyclohexenyl, acenaphthylenyl, and anthracenyl,  
biphenyl, bibenzyl and related bibenzyl homologues  
represented by the formula (a);


20

-21-




where n is a number from 1 to 8.

Particularly preferred are compounds wherein for R<sub>1</sub>  
 5 the combined group -(L<sub>1</sub>)-R<sub>11</sub> is selected from the group  
 consisting of




or



where R<sub>12</sub> is a radical independently selected from halo,  
 C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, -S-(C<sub>1</sub>-C<sub>8</sub> alkyl), -O-(C<sub>1</sub>-C<sub>8</sub>  
 10 alkyl) and C<sub>1</sub>-C<sub>8</sub> haloalkyl where t is a number from 0 to  
 5 and u is a number from 0 to 4 is the group -(L<sub>1</sub>)-R<sub>11</sub>;  
 where, -(L<sub>1</sub>)- is a divalent linking group of 1 to 8  
 atoms and where R<sub>11</sub> is a group selected from (a) or (b).

15 Preferred for R<sub>11</sub> is -(CH<sub>2</sub>)<sub>m</sub>-R<sup>12</sup> wherein m is an  
 integer from 1 to 6, and R<sup>12</sup> is (d) a group represented by  
 the formula:

-22-



wherein a, c, e, n, q, and t are independently an integer from 0 to 2,  $R^{13}$  and  $R^{14}$  are independently selected from a halogen,  $C_1$  to  $C_8$  alkyl,  $C_1$  to  $C_8$  alkyloxy,  $C_1$  to  $C_8$  alkylthio, aryl, heteroaryl, and  $C_1$  to  $C_8$  haloalkyl,  $\alpha$  is an oxygen atom or a sulfur atom,  $L^5$  is a bond,  $-(CH_2)_v-$ ,  $-C=C-$ ,  $-CC-$ ,  $-O-$ , or  $-S-$ , v is an integer from 0 to 2,  $\beta$  is  $-CH_2-$  or  $-(CH_2)_2-$ ,  $\gamma$  is an oxygen atom or a sulfur atom, b is an integer from 0 to 3, d is an integer from 0 to 4, f, p, and w are independently an integer from 0 to 5, r is an integer from 0 to 7, and u is an integer

-23-

from 0 to 4, or is (e) a member of (d) substituted with at least one substituent selected from the group consisting of C<sub>1</sub> to C<sub>6</sub> alkyl, C<sub>1</sub> to C<sub>8</sub> alkyloxy, C<sub>1</sub> to C<sub>8</sub> haloalkyloxy, C<sub>1</sub> to C<sub>8</sub> haloalkyl, aryl, and a halogen.

5

**Preferred R<sub>2</sub> substituents:**


R<sub>2</sub> is preferably selected from the group consisting of hydrogen, C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, -O-(C<sub>1</sub>-C<sub>3</sub> alkyl),

10 -S-(C<sub>1</sub>-C<sub>3</sub> alkyl), -C<sub>3</sub>-C<sub>4</sub> cycloalkyl -CF<sub>3</sub>, halo, -NO<sub>2</sub>, -CN, -SO<sub>3</sub>. Particularly preferred R<sub>2</sub> groups are selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl, -F, -CF<sub>3</sub>, -Cl, -Br, or -O-CH<sub>3</sub>.

15 **Preferred R<sub>3</sub> substituents:**

A preferred subclass of compounds of formula (I) are those wherein X is oxygen.

Another preferred subclass of compounds of  
20 formula (I) are those wherein Z is an oxime amide group.




-24-

Also preferred are compounds of formula (I) wherein  $R_3$  is an oxime amide group and  $R_a$  is hydrogen, methyl or ethyl. For the group  $R_3$  it is preferred that the linking group  $-(L_3)-$  be a bond.

5

**Preferred  $R_4$  substituents:**

Another preferred subclass of compounds of formula (I) are those wherein  $R_4$  is a substituent having an acylamino acid linker with an acylamino acid linker length of 2 or 3 and the acylamino acid linker group,  $-(L_c)-$ , for  $R_4$  is selected from a group represented by the formula;








where  $Q_2$  is selected from the group  $-(CH_2)-$ ,  $-O-$ ,  $-NH-$ ,  $-C(O)-$ , and  $-S-$ , and each  $R_{40}$  is independently selected from hydrogen,  $C_1$ - $C_8$  alkyl, aryl,  $C_1$ - $C_8$  alkaryl,  $C_1$ - $C_8$  alkoxy, aralkyl, and halo. Most preferred are compounds where the acylamino acid linker,  $-(L_c)-$ , for  $R_4$  is selected from the specific groups;

20




-25-



where R<sub>40</sub> is hydrogen or C<sub>1</sub> -C<sub>8</sub> alkyl.

Preferred as the (acylamino acid group) in the group R<sub>4</sub>  
5 is the group:



-26-





wherein  $R^{4a}$  is selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, heteroaryl and aryl; and wherein  $NR^{4b}$  is an amino acid residue of either a natural or unnatural amino acid with the nitrogen atom being part of the amino group of the amino acid. A preferred  $R^{4a}$  group is the group hydrogen (H). A preferred source of amino acid residue is the amino acid group selected from the group comprising isoleucine, valine, phenylalanine, aspartic acid, leucine, glycine and isomers and derivatives thereof. A salt or a prodrug derivative of the (acylamino acid group) is also a suitable substituent.

Particularly preferred are  $R^{4b}$  groups that combine with the nitrogen atom to represent amino acid residues from the amino acid groups selected from: glycine, glycine methyl ester, L-alanine, L-alanine methylester, L-leucine, L-leucine methyl ester, L-aspartic acid, L-aspartic acid dimethyl ester, L-phenyl alanine, L-phenylalanine methyl ester, malonic acid, malonic acid dimethylester, L-valine, L-valine methyl ester, L-isoleucine, L-isoleucine methyl ester, or salt, and derivatives thereof.





**Preferred  $R_5$  Substituents:**

Preferred acid linker,  $-(L_a)-$ , for  $R_5$  is selected from the group consisting of;


-27-



-28-



and



-29-

wherein R<sub>54</sub>, R<sub>55</sub>, R<sub>56</sub> and R<sub>57</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> haloalkyl, aryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, or halo. Preferred (acidic group) for R<sub>5</sub> is selected from the group consisting of -CO<sub>2</sub>H, -SO<sub>3</sub>H and

5 -P(O)(OH)<sub>2</sub>.

**Preferred R<sub>6</sub> and R<sub>7</sub> substituents:**

Another preferred subclass of compounds of formula (I) are those wherein for R<sub>6</sub> and R<sub>7</sub> the non-

10 interfering substituent is independently methyl, ethyl, propyl, isopropyl, thiomethyl, -O-methyl, C<sub>4</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> aralkyl, C<sub>7</sub>-C<sub>12</sub> alkaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, phenyl, tolulyl, xylenyl, biphenyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub>

15 alkenyloxy, C<sub>2</sub>-C<sub>6</sub> alkynyloxy, C<sub>2</sub>-C<sub>12</sub> alkoxyalkyl, C<sub>2</sub>-C<sub>12</sub> alkoxyalkyloxy, C<sub>2</sub>-C<sub>12</sub> alkylcarbonyl, C<sub>2</sub>-C<sub>12</sub> alkylcarbonylamino, C<sub>2</sub>-C<sub>12</sub> alkoxyamino, C<sub>2</sub>-C<sub>12</sub> alkoxyaminocarbonyl, C<sub>1</sub>-C<sub>12</sub> alkylamino, C<sub>1</sub>-C<sub>6</sub> alkylthio, C<sub>2</sub>-C<sub>12</sub> alkylthiocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>

20 alkylsulfonyl, C<sub>2</sub>-C<sub>6</sub> haloalkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkylsulfonyl, C<sub>2</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl, -C(O)O(C<sub>1</sub>-C<sub>6</sub> alkyl), -(CH<sub>2</sub>)<sub>n</sub>-O-(C<sub>1</sub>-C<sub>6</sub> alkyl), benzyloxy, phenoxy, phenylthio, -(CONHSO<sub>2</sub>R), -CHO, amino, amidino, bromo, carbamyl, carboxyl, carbalkoxy, -(CH<sub>2</sub>)<sub>n</sub>-CO<sub>2</sub>H,

25 chloro, cyano, cyanoguanidiny, fluoro, guanidino, hydrazide, hydrazino, hydrazido, hydroxy, hydroxyamino,


-30-

iodo, nitro, phosphono,  $-\text{SO}_3\text{H}$ , thioacetal, thiocarbonyl, and carbonyl; where  $n$  is from 1 to 8.

Most preferred as non-interfering substituents are  
5 methyl, ethyl, propyl, and isopropyl.

Preferred compounds of the invention are those having the general formula (II), or a pharmaceutically acceptable salt, solvate or prodrug derivative thereof;

10







wherein ;


15  $\text{R}_{22}$  is selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl,  $-\text{F}$ ,  $-\text{CF}_3$ ,  $-\text{Cl}$ ,  $-\text{Br}$ , or  $-\text{O}-\text{CH}_3$ ;

-31-

wherein  $R^{4a}$  is selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, heteroaryl and aryl; and wherein  $NR^{4b}$  is an amino acid residue of either a natural or unnatural amino acid with the nitrogen atom being part of the amino group of the amino acid. A preferred  $R^{4a}$  group is the group hydrogen (H); and - (L<sub>4</sub>)- is a divalent group selected from;



-32-



where R<sub>40</sub>, R<sub>41</sub>, R<sub>42</sub>, and R<sub>43</sub> are each independently selected from hydrogen or C<sub>1</sub>-C<sub>8</sub> alkyl.

R<sub>16</sub> is selected from hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, C<sub>1</sub>-C<sub>8</sub> alkylthio C<sub>1</sub>-C<sub>8</sub> haloalkyl, C<sub>1</sub>-C<sub>8</sub> hydroxyalkyl, and halo.

R<sub>13</sub> is selected from hydrogen and C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, -S-(C<sub>1</sub>-C<sub>8</sub> alkyl), C<sub>1</sub>-C<sub>8</sub> haloalkyl, C<sub>1</sub>-C<sub>8</sub> phenyl, halophenyl, hydroxyalkyl, and halo, and t is an integer from 0 to 5.

Preferred specific compounds (and all pharmaceutically acceptable salts, solvates and prodrug derivatives thereof) which are illustrative of the compounds of the invention are as follow:

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]glycine ;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]glycine methyl ester;



13-06-2001

X-12420

009446733(11-07-2000) US00/16319(11-07-2000)

P:DESCPAMD

-33-

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]glycine;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-alanine;

5 N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-alanine methyl ester;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-alanine;

10 N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-leucine;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-leucine methyl ester;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-leucine;

15 N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-aspartic acid;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-aspartic acid dimethyl ester;

20 N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-aspartic acid;

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-  
1H-indol-4-yl]oxy]acetyl]-L-phenylalanine;

-34-

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-phenylalanine methyl ester;

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-phenylalanine;

5 [2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid;

[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid dimethyl ester

10 [2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid;

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-valine;

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-valine methyl ester;

15 *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-valine;

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine;

20 *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine methyl ester; and

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine.

-35-

The salts of the above indole compounds represented by formulae (I) and (II) are an additional aspect of the invention. In those instances where the compounds of the invention possess acidic or basic functional groups  
5 various salts may be formed which are more water soluble and physiologically suitable than the parent compound. Representative pharmaceutically acceptable salts, include but are not limited to, the alkali and alkaline earth salts such as lithium, sodium, potassium, calcium,  
10 magnesium, aluminum and the like. Salts are conveniently prepared from the free acid by treating the acid in solution with a base or by exposing the acid to an ion exchange resin.

15 Included within the definition of pharmaceutically acceptable salts are the relatively non-toxic, inorganic and organic base addition salts of compounds of the present invention, for example, ammonium, quaternary ammonium, and amine cations, derived from nitrogenous  
20 bases of sufficient basicity to form salts with the compounds of this invention (see, for example, S. M. Berge, et al., "Pharmaceutical Salts," J. Phar. Sci., 66: 1-19 (1977)). Moreover, the basic group(s) of the compound of the invention may be reacted with suitable  
25 organic or inorganic acids to form salts such as acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate,

-36-

bitartrate, borate, bromide, camsylate, carbonate,  
chloride, clavulanate, citrate, chloride, edetate,  
edisylate, estolate, esylate, fluoride, fumarate,  
gluceptate, gluconate, glutamate, glycolylarsanilate,  
5 hexylresorcinate, bromide, chloride, hydroxynaphthoate,  
iodide, isothionate, lactate, lactobionate, laurate,  
malate, malseate, mandelate, mesylate, methylbromide,  
methylnitrate, methylsulfate, mucate, napsylate, nitrate,  
oleate, oxalate, palmitate, pantothenate, phosphate,  
10 polygalacturonate, salicylate, stearate, subacetate,  
succinate, tannate, tartrate, tosylate, trifluoroacetate,  
trifluoromethane sulfonate, and valerate.

Certain compounds of the invention may possess one or  
15 more chiral centers, and may thus exist in optically active  
forms. Likewise, when the compounds contain an alkenyl or  
alkenylene group there exists the possibility of cis- and  
trans- isomeric forms of the compounds. The R- and S-  
isomers and mixtures thereof, including racemic mixtures  
20 as well as mixtures of cis- and trans- isomers, are  
contemplated by this invention. Additional asymmetric  
carbon atoms can be present in a substituent group such as  
an alkyl group. All such isomers as well as the mixtures  
thereof are intended to be included in the invention. If  
25 a particular stereoisomer is desired, it can be prepared  
by methods well known in the art by using stereospecific

-37-

reactions with starting materials which contain the asymmetric centers and are already resolved or, alternatively by methods which lead to mixtures of the stereoisomers and subsequent resolution by known methods.

- 5 For example, a racemic mixture may be reacted with a single enantiomer of some other compound. This changes the racemic form into a mixture of diastereomers and diastereomers, because they have different melting points, different boiling points, and different solubilities can  
10 be separated by conventional means, such as crystallization.

Prodrugs are derivatives of the compounds of the invention which have chemically or metabolically cleavable  
15 groups and become by solvolysis or under physiological conditions the compounds of the invention which are pharmaceutically active in vivo. Derivatives of the compounds of this invention have activity in both their acid and base derivative forms, but the acid derivative  
20 form often offers advantages of solubility, tissue compatibility, or delayed release in a mammalian organism (see, Bundgard, H., Design of Prodrugs, pp. 7-9, 21-24, Elsevier, Amsterdam 1985). Prodrugs include acid derivatives well known to practitioners of the art, such  
25 as, for example, esters prepared by reaction of the parent acidic compound with a suitable alcohol, or amides

-38-

prepared by reaction of the parent acid compound with a suitable amine. Simple aliphatic or aromatic esters derived from acidic groups pendent on the compounds of this invention are preferred prodrugs. In some cases it is desirable to prepare double ester type prodrugs such as (acyloxy) alkyl esters or ((alkoxycarbonyl)oxy)alkyl esters. Particularly preferred esters as prodrugs are methyl, ethyl, propyl, isopropyl, n-butyl, isobutyl, tert-butyl, morpholinoethyl, and N,N-diethylglycolamido.

10

N,N-diethylglycolamido ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula (I) (in a medium such as dimethylformamide) with 2-chloro-N,N-diethylacetamide (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA; Item No. 25,099-6).

Morpholinylethyl ester prodrugs may be prepared by reaction of the sodium salt of a compound of Formula (I) (in a medium such as dimethylformamide) 4-(2-chloroethyl)morpholine hydrochloride (available from Aldrich Chemical Co., Milwaukee, Wisconsin USA, Item No. C4,220-3).


a) The 1H-indole-3-glyoxylamide amino derivative compounds of the invention are prepared by room temperature base catalyzed condensation of the amino acid protected at the acid terminus by protecting group

-39-

known in the literature but preferably as the methyl ester with the 1H-indole-3-glyoxylamide acid derivative compound of formula (1) as shown in Scheme I:

5

Scheme 1



Typically, the condensation or coupling is performed in a solvent such as dimethyl formamide, tetrahydrofuran or aqueous mixtures of the like. In general protic solvents are preferred for the purpose of this invention. The reaction is catalyzed by a base including weak organic or inorganic bases. Organic bases such as collidine are preferred. The reaction is also preferably run in the presence of agents that retard or reduce racemization of the amino acid or its

-40-

derivative, such as for example, benzotriazolyl-N-oxy-  
tris(dimethylamino)phosphonium hexafluorophosphate.

Upon completion of the reaction, the mixture is  
concentrated in vacuo. The resulting product mixture is  
5 chromatographed to obtain the target compound.


One of skill in the art is aware that the derivatives  
of the acid such as the acid salt or the methyl ester of  
the acid, can be reacted with the amino acid or  
10 derivatives thereof to obtain the protected compound 2 or  
a corresponding derivative. Such methods are well known  
in the arts and can be found in reference texts such as  
for example J. March Advanced Organic Chemistry, Wiley  
Interscience publishers, New York, N.Y, 1985, and R. C.  
15 Larock Comprehensive Organic Transformations, VCH  
Publishers, New York, N.Y, 1989. The protected compounds  
of formula (2) are also useful sPLA<sub>2</sub> inhibitors and are  
also compounds of this invention.

20        b)        1H-indole-3-acetamide amino acid derivative  
sPLA<sub>2</sub> inhibitors are similarly prepared by condensation of  
the protected amino acid with the 1H-indole-3-acetamide  
sPLA<sub>2</sub> inhibitor. The 1H-indole-3-acetamide sPLA<sub>2</sub>  
inhibitors and methods of making them are set out in U.S.  
25 Patent No. 5,684,034, the entire disclosure of which is  
incorporated herein by reference. Indole-3-acetamide



-41-

amino acid derivative sPLA2 inhibitors of this invention are represented by compounds of formula (IIb), and pharmaceutically acceptable salts and prodrug derivatives thereof,



5

wherein ;

X is oxygen or sulfur;

R<sub>11</sub> is selected from groups (i), (ii) (iii) and (iv)

10 where;

(i) is C<sub>6</sub>-C<sub>20</sub> alkyl, C<sub>6</sub>-C<sub>20</sub> alkenyl, C<sub>6</sub>-C<sub>20</sub> alkynyl, C<sub>6</sub>-C<sub>20</sub> haloalkyl, C<sub>4</sub>-C<sub>12</sub> cycloalkyl, or


(ii) is aryl or aryl substituted by halo, nitro, -CN, -CHO, -OH, -SH, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkylthio, C<sub>1</sub>-

15 C<sub>10</sub> alkoxy, carboxyl, amino, or hydroxyamino; or

(iii) is -(CH<sub>2</sub>)<sub>n</sub>-(R<sub>80</sub>), or -(NH)-(R<sub>81</sub>), where n is 1 to 8, and R<sub>80</sub> is a group recited in (i), and R<sub>81</sub> is selected from a group recited in (i) or (ii);

-42-

(iv) is



where R<sub>87</sub> is hydrogen or C<sub>1</sub>-C<sub>10</sub> alkyl, and R<sub>88</sub> is selected from the group; phenyl, naphthyl, indenyl, and biphenyl, unsubstituted or substituted by halo, -CN, -CHO, -OH, -SH, C<sub>1</sub>-C<sub>10</sub> alkylthio, C<sub>1</sub>-C<sub>10</sub> alkoxy, phenyl, nitro, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> haloalkyl, carboxyl, amino, hydroxyamino; or a substituted or unsubstituted 5 to 8 membered heterocyclic ring;

R<sub>12</sub> is halo, C<sub>1</sub>-C<sub>2</sub> alkylthio, C<sub>1</sub>-C<sub>2</sub> alkyl, C<sub>1</sub>-C<sub>2</sub> alkyaryl or C<sub>1</sub>-C<sub>2</sub> alkoxy;

each R<sub>13</sub> is independently hydrogen, halo, or methyl;


R<sup>14</sup> is the group -L<sub>C</sub>-[acylamino acid], wherein the acylamino acid group is -C(O)-NR<sup>14a</sup>R<sup>14b</sup> wherein R<sup>14a</sup> is selected from the group consisting of H, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, heteroaryl; and -L<sub>C</sub>- is as defined supra, and wherein NR<sup>14b</sup> is an amino acid residue of either a natural or unnatural amino acid with the nitrogen atom being part of the amino group of the amino acid. Most preferred are compounds of formula II wherein the group R<sup>14a</sup> is a hydrogen atom (H). A preferred source of the amino acid residue NR<sup>14b</sup> is an amino acid selected from the group comprising isoleucine, valine, phenylalanine,

-43-

aspartic acid, leucine, glycine and isomers and derivatives thereof;

R<sub>15</sub> is selected from hydrogen, a non-interfering substituent, or the group, -(L<sub>a</sub>)-(acidic group); wherein  
5 -(L<sub>a</sub>)-, is an acid linker having an acid linker length of 1 to 8;

R<sub>16</sub> and R<sub>17</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkenyl, C<sub>1</sub>-C<sub>10</sub> alkynyl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, aryl, aralkyl, or any two adjacent hydrocarbyl groups in  
10 the set R<sub>15</sub>, R<sub>16</sub>, and R<sub>17</sub>, combine with the ring carbon atoms to which they are attached to form a 5 or 6 membered substituted or unsubstituted carbocyclic ring; or C<sub>1</sub>-C<sub>10</sub> haloalkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, C<sub>1</sub>-C<sub>10</sub> haloalkoxy, C<sub>4</sub>-C<sub>8</sub> cycloalkoxy, phenoxy, halo, hydroxy, carboxyl, -SH, -CN,  
15 C<sub>1</sub>-C<sub>10</sub> alkylthio, arylthio, thioacetal, -C(O)O(C<sub>1</sub>-C<sub>10</sub> alkyl), hydrazide, hydrazino, hydrazido, -NH<sub>2</sub>, -NO<sub>2</sub>, -NR<sub>82</sub>R<sub>83</sub>, and -C(O)NR<sub>82</sub>R<sub>83</sub>, where, R<sub>82</sub> and R<sub>83</sub> are independently hydrogen, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> hydroxyalkyl, or taken together with N, R<sub>82</sub> and R<sub>83</sub> form a 5- to 8-  
20 membered heterocyclic ring; or a group having the formula;



where,

-44-





R<sub>84</sub> and R<sub>85</sub> are each independently selected from hydrogen, C<sub>1</sub>-C<sub>10</sub> alkyl, hydroxy, or R<sub>84</sub> and R<sub>85</sub> taken together are =O;

p is 1 to 5,

5 Z is a bond, -O-, -N(C<sub>1</sub>-C<sub>10</sub> alkyl)-, -NH-, or -S-;


and

Q is -CON(R<sub>82</sub>R<sub>83</sub>), -5-tetrazolyl, -SO<sub>3</sub>H,




10

-45-



where n is 1 to 8, R<sub>86</sub> is independently selected from  
hydrogen, a metal, or C<sub>1</sub>-C<sub>10</sub> alkyl, and R<sub>99</sub> is selected  
5 from hydrogen or C<sub>1</sub>-C<sub>10</sub> alkyl.

c) Indole-3-Oxime amide compounds of the invention  
are represented by compounds of formula (III) or a  
pharmaceutically acceptable salt, solvate or prodrug  
10 thereof;



-46-

wherein ;

$R_1$  is selected from groups (a), (b), and (c)


wherein;

5 (a) is C7-C20 alkyl, C7-C20 haloalkyl, C7-C20 alkenyl, C7-C20 alkynyl, carbocyclic radical, or heterocyclic radical, or

(b) is a member of (a) substituted with one or more independently selected non-interfering  
10 substituents; or

(c) is the group  $-(L_1)-R_{11}$ ; where,  $-(L_1)-$  is a divalent linking group of 1 to 8 atoms and where  $R_{11}$  is a group selected from (a) or (b).

15 Particularly preferred are compounds wherein for  $R_1$  the combined group  $-(L_1)-R_{11}$  is selected from the group consisting of




-47-

where  $R_{12}$  is a radical independently selected from halo,  $C_1$ - $C_8$  alkyl,  $C_1$ - $C_8$  alkoxy,  $-S-(C_1-C_8 \text{ alkyl})$ ,  $-O-(C_1-C_8 \text{ alkyl})$  and  $C_1$ - $C_8$  haloalkyl where  $t$  is a number from 0 to 5 and  $u$  is a number from 0 to 4.

5

Also preferred for  $R_{11}$  is  $-(CH_2)_m-R^{12}$  wherein  $m$  is an integer from 1 to 6, and  $R^{12}$  is (d) a group represented by the formula:



10

wherein  $a$ ,  $c$ ,  $e$ ,  $n$ ,  $q$ , and  $t$  are independently an integer from 0 to 2,  $R^{13}$  and  $R^{14}$  are independently selected from a halogen,  $C_1$  to  $C_8$  alkyl,  $C_1$  to  $C_8$  alkyloxy,  $C_1$  to  $C_8$  alkylthio, aryl, heteroaryl, and  $C_1$  to

-48-

C<sub>8</sub> haloalkyl, α is an oxygen atom or a sulfur atom, L<sup>5</sup> is a bond, -(CH<sub>2</sub>)<sub>v</sub>-,

-C=C-, -CC-, -O-, or -S-, v is an integer from 0 to 2, β

is -CH<sub>2</sub>- or -(CH<sub>2</sub>)<sub>2</sub>-, γ is an oxygen atom or a sulfur

5 atom, b is an integer from 0 to 3, d is an integer from

0 to 4, f, p, and w are independently an integer from 0

to 5, r is an integer from 0 to 7, and u is an integer


from 0 to 4, or is (e) a member of (d) substituted with

at least one substituent selected from the group

10 consisting of C<sub>1</sub> to C<sub>6</sub> alkyl, C<sub>1</sub> to C<sub>8</sub> alkyloxy, C<sub>1</sub> to C<sub>8</sub> haloalkyloxy, C<sub>1</sub> to C<sub>8</sub> haloalkyl, aryl, and a halogen.


R<sub>2</sub> is hydrogen, or a group containing 1 to 4 non-hydrogen atoms plus any required hydrogen atoms;

15 -(L<sub>3</sub>)- Z, is the group where -(L<sub>3</sub>)- is a divalent linker group selected from a bond or a divalent group selected from:





and Z is selected from an oxime amide or oxime thioamide group represented by the formulae,



5

wherein, X is oxygen or sulfur; and R<sub>a</sub> is selected from hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, aryl, C<sub>1</sub>-C<sub>8</sub> alkaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, aralkyl and -CN;

10 R<sub>4</sub> is the group, -(L<sub>C</sub>)-(acylamino acid group); wherein -(L<sub>C</sub>)-, is an acylamino acid linker having an acylamino acid linker length of 1 to 8;


R<sub>5</sub> is selected from hydrogen, a non-interfering substituent, or the group, -(L<sub>a</sub>)-(acidic group); wherein  
15 -(L<sub>a</sub>)-, is an acid linker having an acid linker length of 1 to 8.

R<sub>6</sub> and R<sub>7</sub> are selected from hydrogen, non-interfering substituent, carbocyclic radical,  
20 carbocyclic radical substituted with non-interfering substituent(s), heterocyclic radicals, and heterocyclic radical substituted with non-interfering substituent(s).

-50-


**Preferred Subgroups of Compounds of Formula (III):**  
**Preferred R<sub>1</sub> substituents:**

A preferred subclass of compounds of formula (III)  
5 are those where for R<sub>1</sub> the divalent linking group -(L<sub>1</sub>)-  
is a group represented by any one of the following  
formulae (Ia), (Ib), (Ic), (Id), (Ie), or (If):



10

or




15 where Q<sub>1</sub> is a bond or any of the divalent groups (Ia),  
(Ib), (Ic), (Id), (Ie), and (If) and each R<sub>10</sub> is

-51-

independently hydrogen, C<sub>1-8</sub> alkyl, C<sub>1-8</sub> haloalkyl or C<sub>1-8</sub> alkoxy.

Particularly preferred as the linking group -(L<sub>1</sub>)- of  
5 R<sub>1</sub> is an alkylene chain of 1 or 2 carbon atoms, namely,  
-(CH<sub>2</sub>)- or -(CH<sub>2</sub>-CH<sub>2</sub>)-.


The preferred group for R<sub>11</sub> is a substituted or  
unsubstituted group selected from the group consisting of  
10 C<sub>5</sub>-C<sub>14</sub> cycloalkyl, C<sub>5</sub>-C<sub>14</sub> cycloalkenyl, phenyl, naphthyl,  
norbornanyl, bicycloheptadienyl, tolulyl, xylenyl,  
indenyl, stilbenyl, terphenyl, diphenylethylenyl,  
phenyl-cyclohexenyl, acenaphthylenyl, and anthracenyl,  
biphenyl, bibenzylyl and related bibenzylyl homologues  
15 represented by the formula (a);




where n is a number from 1 to 8.

20 Particularly preferred are compounds wherein for R<sub>1</sub>  
the combined group -(L<sub>1</sub>)-R<sub>11</sub> is selected from the group  
consisting of

-52-




or



where R<sub>12</sub> is a radical independently selected from halo, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> alkoxy, -S-(C<sub>1</sub>-C<sub>8</sub> alkyl), -O-(C<sub>1</sub>-C<sub>8</sub> alkyl) and C<sub>1</sub>-C<sub>8</sub> haloalkyl where t is a number from 0 to 5 and u is a number from 0 to 4.

Also preferred for  $R_{11}$  is  $-(CH_2)_m-R^{12}$  wherein  $m$  is an integer from 1 to 6, and  $R^{12}$  is (d) a group represented by

10 the formula:



-53-

wherein a, c, e, n, q, and t are independently an integer from 0 to 2,  $R^{13}$  and  $R^{14}$  are independently selected from a halogen,  $C_1$  to  $C_8$  alkyl,  $C_1$  to  $C_8$  alkyloxy,  $C_1$  to  $C_8$  alkylthio, aryl, heteroaryl, and  $C_1$  to  $C_8$  haloalkyl,  $\alpha$  is an oxygen atom or a sulfur atom,  $L^5$  is a bond,  $-(CH_2)_v-$ ,  $-C=C-$ ,  $-CC-$ ,  $-O-$ , or  $-S-$ , v is an integer from 0 to 2,  $\beta$  is  $-CH_2-$  or  $-(CH_2)_2-$ ,  $\gamma$  is an oxygen atom or a sulfur atom, b is an integer from 0 to 3, d is an integer from 0 to 4, f, p, and w are independently an integer from 0 to 5, r is an integer from 0 to 7, and u is an integer from 0 to 4, or is (e) a member of (d) substituted with at least one substituent selected from the group consisting of  $C_1$  to  $C_6$  alkyl,  $C_1$  to  $C_8$  alkyloxy,  $C_1$  to  $C_8$  haloalkyloxy,  $C_1$  to  $C_8$  haloalkyl, aryl, and a halogen.


**Preferred  $R_2$  substituents:**

$R_2$  is preferably selected from the group consisting of hydrogen,  $C_1$ - $C_4$  alkyl,  $C_2$ - $C_4$  alkenyl,  $-O-(C_1$ - $C_3$  alkyl),  $-S-(C_1$ - $C_3$  alkyl),  $-C_3$ - $C_4$  cycloalkyl  $-CF_3$ , halo,  $-NO_2$ ,  $-CN$ ,  $-SO_3$ . Particularly preferred  $R_2$  groups are selected from hydrogen, methyl, ethyl, propyl, isopropyl, cyclopropyl,  $-F$ ,  $-CF_3$ ,  $-Cl$ ,  $-Br$ , or  $-O-CH_3$ .

-54-

**Preferred R<sub>4</sub> substituents:**






Another preferred subclass of compounds of formula (III) are those wherein R<sub>4</sub> is a substituent having an acylamino acid linker with an acylamino acid linker length of 2 or 3 and the acylamino acid linker group, -(L<sub>C</sub>)-, for R<sub>4</sub> is selected from a group represented by the formula;



10

where Q<sub>2</sub> is selected from the group -(CH<sub>2</sub>)-, -O-, -NH-, -C(O)-, and -S-, and each R<sub>40</sub> is independently selected from hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, aryl, C<sub>1</sub>-C<sub>8</sub> alkaryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, aralkyl, and halo. Most preferred are compounds where the acylamino acid linker, -(L<sub>C</sub>)-, for R<sub>4</sub> is selected from the specific groups;


-55-



where R<sub>40</sub> is hydrogen or C<sub>1</sub> -C<sub>8</sub> alkyl.

Preferred as the (acylamino acid group) in the group R<sub>4</sub>

5 is the group:



-56-

wherein R<sup>4a</sup> is selected from the group consisting of H,  
(C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, heteroaryl and aryl; and  
wherein NR<sup>4b</sup> is an amino acid residue of either a natural  
or unnatural amino acid with the nitrogen atom being part  
5 of the amino group of the amino acid. A preferred R<sup>4a</sup>  
group is the group hydrogen (H). A preferred source of  
amino acid residue is the amino acid group selected from  
the group comprising isoleucine, valine, phenylalanine,  
aspartic acid, leucine, glycine and isomers and  
10 derivatives thereof.

A salt or a prodrug derivative of the (acylamino  
acid group) is also a suitable substituent.

15 Particularly preferred are R<sup>4b</sup> groups that combine  
with the nitrogen atom to represent amino acid groups  
selected from: glycine, glycine methyl ester, L-  
alanine, L-alanine methylester, L-leucine, L-leucine  
methyl ester, L-aspartic acid, L-aspartic acid dimethyl  
20 ester, L-phenyl alanine, L-phenylalanine methyl ester,  
malonic acid, malonic acid dimethylester, L- valine, L-  
valine methyl ester, L-isoleucine, L-isoleucine methyl  
ester, or salt, and derivatives thereof.





-57-






**Preferred R<sub>5</sub> Substituents:**

Preferred acid linker,  $-(L_a)-$ , for R<sub>5</sub> is selected from the group consisting of;


5




-58-



-59-



and



wherein R<sub>54</sub>, R<sub>55</sub>, R<sub>56</sub> and R<sub>57</sub> are each independently hydrogen, C<sub>1</sub>-C<sub>8</sub> alkyl, C<sub>1</sub>-C<sub>8</sub> haloalkyl, aryl, C<sub>1</sub>-C<sub>8</sub> alkoxy, or halo. Preferred (acidic group) for R<sub>5</sub> is selected from the group consisting of -CO<sub>2</sub>H, -SO<sub>3</sub>H and -P(O)(OH)<sub>2</sub>

**Preferred R<sub>6</sub> and R<sub>7</sub> substituents:**

Another preferred subclass of compounds of formula (III) are those wherein for R<sub>6</sub> and R<sub>7</sub> the non-interfering substituent is independently methyl, ethyl, propyl, isopropyl, thiomethyl, -O-methyl, C<sub>4</sub>-C<sub>6</sub> alkyl, C<sub>2</sub>-C<sub>6</sub> alkenyl, C<sub>2</sub>-C<sub>6</sub> alkynyl, C<sub>7</sub>-C<sub>12</sub> aralkyl, C<sub>7</sub>-C<sub>12</sub> alkaryl, C<sub>3</sub>-C<sub>8</sub> cycloalkyl, C<sub>3</sub>-C<sub>8</sub> cycloalkenyl, phenyl, tolulyl, xylenyl, biphenyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>2</sub>-C<sub>6</sub> alkenyloxy, C<sub>2</sub>-C<sub>6</sub> alkynyloxy, C<sub>2</sub>-C<sub>12</sub> alkoxyalkyl, C<sub>2</sub>-C<sub>12</sub>

-60-

alkoxyalkyloxy, C<sub>2</sub>-C<sub>12</sub> alkylcarbonyl, C<sub>2</sub>-C<sub>12</sub>

alkylcarbonylamino, C<sub>2</sub>-C<sub>12</sub> alkoxyamino, C<sub>2</sub>-C<sub>12</sub>

alkoxyaminocarbonyl, C<sub>1</sub>-C<sub>12</sub> alkylamino, C<sub>1</sub>-C<sub>6</sub> alkylthio,

C<sub>2</sub>-C<sub>12</sub> alkylthiocarbonyl, C<sub>1</sub>-C<sub>6</sub> alkylsulfinyl, C<sub>1</sub>-C<sub>6</sub>

5 alkylsulfonyl, C<sub>2</sub>-C<sub>6</sub> haloalkoxy, C<sub>1</sub>-C<sub>6</sub>

haloalkylsulfonyl, C<sub>2</sub>-C<sub>6</sub> haloalkyl, C<sub>1</sub>-C<sub>6</sub> hydroxyalkyl,

-C(O)O(C<sub>1</sub>-C<sub>6</sub> alkyl), -(CH<sub>2</sub>)<sub>n</sub>-O-(C<sub>1</sub>-C<sub>6</sub> alkyl), benzyloxy,

phenoxy, phenylthio, -(CONHSO<sub>2</sub>R), -CHO, amino, amidino,

bromo, carbamyl, carboxyl, carbalkoxy, -(CH<sub>2</sub>)<sub>n</sub>-CO<sub>2</sub>H,

10 chloro, cyano, cyanoguanidinyl, fluoro, guanidino,

hydrazide, hydrazino, hydrazido, hydroxy, hydroxyamino,

iodo, nitro, phosphono, -SO<sub>3</sub>H, thioacetal, thiocarbonyl,

and carbonyl; where n is from 1 to 8.


15 Most preferred as non-interfering substituents are  
methyl, ethyl, propyl, and isopropyl.

The indole-3-oxime compounds of the invention can be  
prepared following protocol of scheme 2 below;

20

-61-

Scheme 2



5

To introduce the oxime functionality, the methyl ester of the glyoxylamide (compound 10 in scheme 1, compound 1 in scheme 2, *supra.*) is heated with hydroxylamine hydrochloride (when R is H) in a THF/methanol mixture for 8 hours or until the reaction was deemed complete. The reaction product is isolated by chromatography or other known laboratory procedure to afford a white solid. Substituted oximes such as when R is methyl, ethyl, phenyl or other substituent can be prepared by reacting the corresponding substituted hydroxylamine hydrochloride or free base with the

-62-

glyoxylamide as described supra. The ester functionality at the 4 or 5 position on the indole nucleus, as in for example, compound 2, can be: (a) converted to the acid by hydrolysis using lithium hydroxide or other known ester hydrolysis methods to afford compounds of formula 3, or (b) converted to an amide functionality directly or via the acid functionality to afford compounds of formula 4. General procedures for the conversion of organic acids to amino acid are well known to artisans in the field, and have been documented in general reference texts including, for example, J. March Advanced Organic Chemistry, Wiley Interscience publishers, New York, N.Y, 1985, and R. C. Larock Comprehensive Organic Transformations, VCH Publishers, New York, N.Y, 1989.

15

The oxime acid compounds of formula 3 such as the methyloxime compound such as that of formula 4 can be converted to the corresponding amino acid derivative via the methylester by coupling with various amino acids by general coupling procedures known to one skilled in the art. Additional references, or procedures are found in J. March Advanced Organic Chemistry, Wiley Interscience publishers, New York, N.Y, 1985; R. C. Larock Comprehensive Organic Transformations, VCH Publishers, New York, N.Y, 1989 and J. Jones Amino Acids and Peptide

-63-


Synthesis, Oxford Science Publications, Stephen G. Davis,  
Editor, Oxford University Press Inc., New York, NY, 1992.

5    **III. Method of Making the 1H-Indole-3-Glyoxylamide  
Starting Material for Preparing the Compounds of the  
Invention:**

10            The synthesis of the indole compounds of the  
invention (viz., Compounds of Formulae I and II) can be  
accomplished by well known methods as recorded in the  
chemical literature. In particular, the indole starting  
materials may be prepared by the synthesis schemes  
taught in US Patent No. 5,654,326; the disclosure of  
15    which is incorporated herein by reference. Another  
method of making 1H-indole-3-glyoxylamide SPLA<sub>2</sub>  
inhibitors is described in United States Patent  
Application Serial No. 09/105381, filed June 26, 1998  
and titled, "Process for Preparing 4-substituted 1-H-  
20    Indole-3-glyoxylamides" the entire disclosure of which is  
incorporated herein by reference.


United States Patent Application Serial  
No. 09/105381 discloses the following process having  
25    steps (a) thru (i):  
Preparing a compound of the formula (Iz) or a  
pharmaceutically acceptable salt or prodrug derivative  
thereof

- 64 -



5 wherein:

$R^{1z}$  is selected from the group consisting of -C<sub>7</sub>-C<sub>20</sub> alkyl,



where

10  $R^{10z}$  is selected from the group consisting of halo, C<sub>1</sub>-C<sub>10</sub> alkyl, C<sub>1</sub>-C<sub>10</sub> alkoxy, -S-(C<sub>1</sub>-C<sub>10</sub> alkyl) and halo(C<sub>1</sub>-C<sub>10</sub>)alkyl, and tz is an integer from 0 to 5 both inclusive;

15  $R^{2z}$  is selected from the group consisting of hydrogen, halo, C<sub>1</sub>-C<sub>3</sub> alkyl, C<sub>3</sub>-C<sub>4</sub> cycloalkyl, C<sub>3</sub>-C<sub>4</sub>



-65-


cycloalkenyl, -O-(C<sub>1</sub>-C<sub>2</sub> alkyl), -S-(C<sub>1</sub>-C<sub>2</sub> alkyl), aryl, aryloxy and HET;

R<sup>4z</sup> is the group -CO<sub>2</sub>H, or salt and prodrug derivative thereof; and

5 R<sup>5z</sup>, R<sup>6z</sup> and R<sup>7z</sup> are each independently selected from the group consisting of hydrogen, (C<sub>1</sub>-C<sub>6</sub>)alkyl, (C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy, halo(C<sub>2</sub>-C<sub>6</sub>)alkyl, bromo, chloro, fluoro, iodo and aryl;

which process comprises the steps of:

10 a) halogenating a compound of formula Xz




Xz

where R<sup>8z</sup> is (C<sub>1</sub>-C<sub>6</sub>)alkyl, aryl or HET;

with SO<sub>2</sub>Cl<sub>2</sub> to form a compound of formula


15 IX




b) hydrolyzing and decarboxylating a compound of formula IXz

20

-66-




to form a compound of formula VIIIz




5

c) alkylating a compound of formula VIIz




10

with a compound of formula VIIIz




to form a compound of formula VIz



15

-67-

- d) aminating and dehydrating a compound of formula VIz




5

with an amine of the formula  $R^{1z}NH_2$  in the presence of a solvent that forms an azeotrope with water to form a compound of formula Vz;

10

- e) oxidizing a compound of formula Vz



15


by refluxing in a polar hydrocarbon solvent having a boiling point of at least 150 °C and a dielectric constant of at least 10 in the presence of a catalyst to form a compound of formula IVz

13-06-2001  
X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)


P DESCAMD

-68-




-69-


f) alkylating a compound of the formula IVz



with an alkylating agent of the formula  $XCH_2R^{4az}$   
 where X is a leaving group and  $R^{4az}$  is  $-CO_2R^{4b}$ ,  
 where  $R^{4bz}$  is an acid protecting group to form a  
 compound of formula IIIz




g) reacting a compound of formula IIIz



with oxalyl chloride and ammonia to form a  
 compound of formula IIz


-70-



IIz; and

5

- h) optionally hydrolyzing a compound of  
formula IIz



10


IIz

to form a compound of formula Iz.

An alternative protocol useful for the synthesis of  
the starting material is shown in Scheme 1 below:

-71-

Scheme 1



5 The synthesis of indole-3-oxime amides (compound of formula I and II, supra.) of this invention uses

-72-

as starting material the glyoxamide ((3-(2-amino-1,2-dioxoethyl)-2-methyl-1-(phenylmethyl)-1H-indol-4-yl)oxy)acetic acid methyl ester, compound 10, *supra*. This starting material is prepared as set out in the preceding  
5 section or by the method of Example 9 of U.S. Patent No. 5,654,326 (the disclosure of which is incorporated herein by reference).

To obtain the glyoxylamide starting material  
10 substituted in the 4-position with an (acidic group) linked through an oxygen atom, the reactions outlined in the scheme *supra*, are used (for conversions 1 through 5, see ref. Robin D. Clark, Joseph M. Muchowski, Lawrence E. Fisher, Lee A. Flippin, David B. Repke, Michel Souchet,  
15 *Synthesis*, 1991, 871-878, the disclosures of which are incorporated herein by reference). The starting material ortho-nitrotoluene, 1, is readily reduced to 2-methyl,3-methoxyaniline, 2. Reduction of 1 is by the catalytic hydrogenation of the corresponding nitrotoluene using  
20 palladium on carbon as catalyst. The reduction can be carried out in ethanol or tetrahydrofuran (THF) or a combination of both, using a low pressure of hydrogen. The aniline 2, obtained, is converted to the N-tert-butyloxycarbonyl derivative 3, in good yield, on heating  
25 with di-tert-butyl dicarbonate in THF at reflux temperature. The dilithium salt of the dianion of 3 is



-73-

generated at -40 to -20°C in THF using sec-butyllithium and reacted with the appropriately substituted N-methoxy-N-methylalkanamide to form the ketone 4. This product (4) may be purified by crystallization from hexane, or reacted  
5 directly with trifluoroacetic acid in methylene chloride to give the 1,3-unsubstituted indole 5. The 1,3-unsubstituted indole 5 is reacted with sodium hydride in dimethylformamide at room temperature (20-25°C) for 0.5-1.0 hour. The resulting sodium salt of 5 is treated with  
10 an equivalent of arylmethyl halide and the mixture stirred at a temperature range of 0-100°C, usually at ambient room temperature, for a period of 4 to 36 hours to give the 1-arylmethylindole, 6. This indole, 6, is O-demethylated by stirring with boron tribromide in methylene chloride for  
15 approximately 5 hours (see ref. Tsung-Ying Shem and Charles A Winter, *Adv. Drug Res.*, 1977, 12, 176, the disclosure of which is incorporated herein by reference). The 4-hydroxyindole, 7, is alkylated with an alpha bromoalkanoic acid ester in dimethylformamide (DMF) using  
20 sodiumhydride as a base, with reaction condition of 5 to 6. The  $\alpha$ -[(indol-4-yl)oxy]alkanoic acid ester, 8, is reacted with oxalyl chloride in methylene chloride to give 9, which is not purified but reacted directly with ammonia to give the glyoxamide 10.

-74-

Glyoxamide starting material compounds substituted at the 5 position of the indole nucleus with an (acidic group) may be prepared by methods and starting materials shown in schemes 2 and 3 of Patent No. 5,654,326; the disclosure of which is incorporated herein by reference.

#### IV. Methods of Using the Compounds of the Invention:

The indole compounds described herein are believed to achieve their beneficial therapeutic action principally by direct inhibition of mammalian (including human) sPLA<sub>2</sub>, and not by acting as antagonists for arachidonic acid, nor other active agents below arachidonic acid in the arachidonic acid cascade, such as 5-lipoxygenases, cyclooxygenases, and etc.

The method of the invention for inhibiting sPLA<sub>2</sub> mediated release of fatty acids comprises contacting mammalian sPLA<sub>2</sub> with an therapeutically effective amount of indole compounds corresponding to Formulae (I) or (II) as described herein including salt or a prodrug derivative thereof.

Another aspect of this invention is a method for treating Inflammatory Diseases such as inflammatory bowel disease, septic shock, adult respiratory distress

-75-

syndrome, pancreatitis, trauma, bronchial asthma,  
allergic rhinitis, rheumatoid arthritis, osteoarthritis,  
and related diseases which comprises administering to a  
mammal (including a human) a therapeutically effective  
5 dose of the indole compound of the invention (see,  
formulae I and II).

As previously noted the compounds of this invention  
are useful for inhibiting sPLA<sub>2</sub> mediated release of  
10 fatty acids such as arachidonic acid. By the term,  
"inhibiting" is meant the prevention or therapeutically  
significant reduction in release of sPLA<sub>2</sub> initiated  
fatty acids by the compounds of the invention. By  
"pharmaceutically acceptable" it is meant the carrier,  
15 diluent or excipient must be compatible with the other  
ingredients of the formulation and not deleterious to  
the recipient thereof.

The specific dose of a compound administered  
20 according to this invention to obtain therapeutic or  
prophylactic effects will, of course, be determined by the  
particular circumstances surrounding the case, including,  
for example, the compound administered, the route of  
administration and the condition being treated. Typical  
25 daily doses will contain a non-toxic dosage level of from

-76-

about 0.01 mg/kg to about 50 mg/kg of body weight of an active compound of this invention.

Preferably compounds of the invention (per Formula I  
5 or II) or pharmaceutical formulations containing these compounds are in unit dosage form for administration to a mammal. The unit dosage form can be a capsule or tablet itself, or the appropriate number of any of these. The quantity of Active ingredient in a unit dose of  
10 composition may be varied or adjusted from about 0.1 to about 1000 milligrams or more according to the particular treatment involved. It may be appreciated that it may be necessary to make routine variations to the dosage depending on the age and condition of the patient. The  
15 dosage will also depend on the route of administration.

The compound can be administered by a variety of routes including oral, aerosol, rectal, transdermal, subcutaneous, intravenous, intramuscular, and  
20 intranasal.

Pharmaceutical formulations of the invention are prepared by combining (e.g., mixing) a therapeutically effective amount of the indole compound of the invention  
25 together with a pharmaceutically acceptable carrier or diluent therefor. The present pharmaceutical

-77-

formulations are prepared by known procedures using well known and readily available ingredients.

In making the compositions of the present invention, the Active ingredient will usually be admixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a capsule, sachet, paper or other container. When the carrier serves as a diluent, it may be a solid, semi-solid or liquid material which acts as a vehicle, or can be in the form of tablets, pills, powders, lozenges, elixirs, suspensions, emulsions, solutions, syrups, aerosols (as a solid or in a liquid medium), or ointment, containing, for example, up to 10% by weight of the active compound. The compounds of the present invention are preferably formulated prior to administration.

For the pharmaceutical formulations any suitable carrier known in the art can be used. In such a formulation, the carrier may be a solid, liquid, or mixture of a solid and a liquid. For example, for intravenous injection the compounds of the invention may be dissolved in at a concentration of 2 mg/ml in a 4% dextrose/0.5% Na citrate aqueous solution. Solid form formulations include powders, tablets and capsules. A solid carrier can be one or more substances which may also

-78-

act as flavoring agents, lubricants, solubilisers, suspending agents, binders, tablet disintegrating agents and encapsulating material.

5           Tablets for oral administration may contain suitable excipients such as calcium carbonate, sodium carbonate, lactose, calcium phosphate, together with disintegrating agents, such as maize, starch, or alginic acid, and/or binding agents, for example, gelatin or  
10   acacia, and lubricating agents such as magnesium stearate, stearic acid, or talc.

In powders the carrier is a finely divided solid which is in admixture with the finely divided Active  
15   ingredient. In tablets the Active ingredient is mixed with a carrier having the necessary binding properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain from about 1 to about 99 weight percent of the Active  
20   ingredient which is the novel compound of this invention. Suitable solid carriers are magnesium carbonate, magnesium stearate, talc, sugar lactose, pectin, dextrin, starch, gelatin, tragacanth, methyl cellulose, sodium carboxymethyl cellulose, low melting waxes, and cocoa  
25   butter.

-79-

Sterile liquid form formulations include suspensions, emulsions, syrups and elixirs.

The Active ingredient can be dissolved or suspended  
5 in a pharmaceutically acceptable carrier, such as sterile  
water, sterile organic solvent or a mixture of both. The  
Active ingredient can often be dissolved in a suitable  
organic solvent, for instance aqueous propylene glycol.  
Other compositions can be made by dispersing the finely  
10 divided Active ingredient in aqueous starch or sodium  
carboxymethyl cellulose solution or in a suitable oil.

The following pharmaceutical formulations 1 thru 8  
are illustrative only and are not intended to limit the  
15 scope of the invention in any way. "Active ingredient",  
refers to a compound according to Formula (I) or (II) or a  
pharmaceutically acceptable salt, solvate, or prodrug  
thereof.

#### Formulation 1

20 Hard gelatin capsules are prepared using the  
following ingredients:

	Quantity (mg/capsule)
Active ingredient	250
Starch, dried	200
Magnesium stearate	<u>10</u>
Total	460 mg

13-06-2001

X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)

P DESCAMD

-80-

### Formulation 2

A tablet is prepared using the ingredients below:

	Quantity (mg/tablet)
Active ingredient	250
Cellulose, microcrystalline	400
Silicon dioxide, fumed	10
Stearic acid	<u>5</u>
Total	665 mg

5

The components are blended and compressed to form tablets each weighing 665 mg

### Formulation 3

10 An aerosol solution is prepared containing the following components:

	Weight
Active ingredient	<u>0.25</u>
Ethanol	25.75
Propellant 22 (Chlorodifluoromethane)	<u>74.00</u>
Total	100.00

15 The active compound is mixed with ethanol and the mixture added to a portion of the propellant 22, cooled to -30°C and transferred to a filling device. The required amount is then fed to a stainless steel container and



13-06-2001

X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)

P DESC PAMD

-81-

diluted with the remainder of the propellant. The valve units are then fitted to the container.

#### Formulation 4

- 5        Tablets, each containing 60 mg of Active ingredient, are made as follows:

Active ingredient	60 mg
Starch	45 mg
Microcrystalline cellulose	35 mg
Polyvinylpyrrolidone (as 10% solution in water)	4 mg
Sodium carboxymethyl starch	4.5 mg
Magnesium stearate	0.5 mg
Talc	<u>1 mg</u>
Total	150 mg

- 10        The Active ingredient, starch and cellulose are passed through a No. 45 mesh U.S. sieve and mixed thoroughly. The aqueous solution containing polyvinylpyrrolidone is mixed with the resultant powder, and the mixture then is passed through a No. 14 mesh U.S. sieve. The granules so produced are dried at 50°C and
- 15        passed through a No. 18 mesh U.S. sieve. The sodium carboxymethyl starch, magnesium stearate and talc, previously passed through a No. 60 mesh U.S. sieve, are then added to the granules which, after mixing, are compressed on a tablet machine to yield tablets each
- 20        weighing 150 mg.

-82-

**Formulation 5**

Capsules, each containing 80 mg of Active ingredient,  
are made as follows:

5

Active ingredient	80 mg
Starch	59 mg
Microcrystalline cellulose	59 mg
Magnesium stearate	<u>2 mg</u>
Total	200 mg

The Active ingredient, cellulose, starch, and  
magnesium stearate are blended, passed through a No. 45  
mesh U.S. sieve, and filled into hard gelatin capsules in  
200 mg quantities.

**Formulation 6**

Suppositories, each containing 225 mg of Active  
ingredient, are made as follows:

Active ingredient	225 mg
Saturated fatty acid glycerides	<u>2,000 mg</u>
Total	2,225 mg

15

The Active ingredient is passed through a No. 60 mesh  
U.S. sieve and suspended in the saturated fatty acid  
glycerides previously melted using the minimum heat  
necessary. The mixture is then poured into a suppository  
mold of nominal 2 g capacity and allowed to cool.

13-06-2001

X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)

P. DESCPAMD

-83-

### Formulation 7

Suspensions, each containing 50 mg of Active ingredient per 5 ml dose, are made as follows:

5

Active ingredient	50 mg
Sodium carboxymethyl cellulose	50 mg
Syrup	1.25 ml
Benzoic acid solution	0.10 ml
Flavor	q.v.
Color	q.v.
Purified water to total	5 ml

The Active ingredient is passed through a No. 45 mesh U.S. sieve and mixed with the sodium carboxymethyl cellulose and syrup to form a smooth paste. The benzoic acid solution, flavor and color are diluted with a portion of the water and added, with stirring. Sufficient water is then added to produce the required volume.

### Formulation 8

15 An intravenous formulation may be prepared as follows:

Active ingredient	100 mg
Isotonic saline	1,000 ml

-84-


The solution of the above ingredients generally is administered intravenously to a subject at a rate of 1 ml per minute.

5 All of the products of the Examples described below as well as intermediates used in the following procedures showed satisfactory nmr and IR spectra. They also had the correct mass spectral values.

10

Example 1

Preparation of [[3-(2-Amino-1,2-dioxoethyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester, a compound represented by the compound of formula (1) formula:



15

Part A. Preparation of 2-Ethyl-4-methoxy-1H-indole.

A solution of 140 mL (0.18 mol) of 1.3M sec-butyl lithium in cyclohexane was added slowly to N-tert-  
20 butoxycarbonyl-3-methoxy-2-methylaniline (21.3g, 0.09 mol)

-85-

in 250 mL of THF keeping the temperature below -40°C with a dry ice-ethanol bath. The bath was removed and the temperature allowed to rise to 0°C and then the bath replaced. After the temperature had cooled to -60°C,

5 18.5g (0.18 mol) of N-methoxy-N-methylpropanamide in an equal volume of THF was added dropwise. The reaction mixture was stirred 5 minutes, the cooling bath removed and stirred an additional 18 hours. It was then poured into a mixture of 300 mL of ether and 400 mL of 0.5N HCl.

10 The organic layer was separated, washed with water, brine, dried over  $\text{MgSO}_4$ , and concentrated at reduced pressure to give 25.5g of a crude of 1-[2-(tert-butoxycarbonylamino)-6-methoxyphenyl]-2-butanone. This material was dissolved in 250 mL of methylene chloride and 50 mL of

15 trifluoroacetic acid and stirred for a total of 17 hours. The mixture was concentrated at reduced pressure and ethyl acetate and water added to the remaining oil. The ethyl acetate was separated, washed with brine, dried ( $\text{MgSO}_4$ ) and concentrated. The residue was chromatographed three

20 times on silica eluting with 20% EtOAc/hexane to give 13.9g of 2-ethyl-4-methoxy-1H-indole.

Analyses for  $\text{C}_{11}\text{H}_{13}\text{NO}$ :

Calculated: C, 75.40; H, 7.48; N, 7.99

Found: C, 74.41; H, 7.64; N, 7.97.

25

**Part B. Preparation of 2-Ethyl-4-methoxy-1-(phenylmethyl)-1H-indole.**

-86-

2-Ethyl-4-methoxy-1H-indole (4.2g, 24 mmol) was dissolved in 30 mL of DMF and 960mg (24 mmol) of 60% NaH/mineral oil was added. After 1.5 hours, 2.9 mL (24 mmol) of benzyl bromide was added. After 4 hours, the mixture was diluted with water and extracted twice with ethyl acetate. The combined ethyl acetate was washed with brine, dried (MgSO<sub>4</sub>) and concentrated at reduced pressure. The residue was chromatographed on silica gel and eluted with 20% EtOAc/hexane to give 3.1g (49% yield) of 2-ethyl-4-methoxy-1-(phenylmethyl)-1H-indole.

**Part C. Preparation of 2-Ethyl-4-hydroxy-1-(phenylmethyl)-1H-indole.**

3.1g (11.7 mmol) of 2-ethyl-4-methoxy-1-(phenylmethyl)-1H-indole was O-demethylated by treating it with 48.6 mL of 1M BBr<sub>3</sub> in methylene chloride with stirring at room temperature for 5 hours, followed by concentration at reduced pressure. The residue was dissolved in ethyl acetate, washed with brine and dried (MgSO<sub>4</sub>). After concentrating at reduced pressure, the residue was chromatographed on silica gel eluting with 20% EtOAc/hexane to give 1.58g (54% yield) of 2-ethyl-4-hydroxy-1-(phenylmethyl)-1H-indole, mp, 86-90°C.

Analyses for C<sub>17</sub>H<sub>17</sub>NO:

Calculated: C, 81.24; H, 6.82; N, 5.57

Found: C, 81.08; H, 6.92; N, 5.41.

**Part D. Preparation of [[2-Ethyl-1-(phenylmethyl)-**

-87-

**1H-indol-4-yl]oxy]acetic acid methyl ester.**

2-ethyl-4-hydroxy-1-(phenylmethyl)-1H-indole (1.56g, 6.2 mmol) was added to a mixture of 248mg (6.2 mmol) of 60% NaH/mineral oil in 20mL DMF and stirred for 0.67 hour.

5

Then 0.6 mL(6.2 mmol) of methyl bromoacetate was added and stirring was continued for 17 hours. The mixture was diluted with water and extracted with ethyl acetate. The ethyl acetate solution was washed with brine, dried (MgSO<sub>4</sub>), and concentrated at reduced pressure. The residue was chromatographed on silica gel eluting with 20% EtOAc/hexane, to give 1.37g (69% yield) of [[2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester, 89-92°C.

15 Analyses for C<sub>20</sub>H<sub>21</sub>NO<sub>3</sub>:

Calculated: C, 74.28; H, 6.55; N, 4.33

Found: C, 74.03; H, 6.49; N, 4.60.

**Part E. Preparation of [[3-(2-Amino-1,2-dioxoethyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester.**

20 Oxalyl chloride (0.4 mL, 4.2 mmol) was added to 1.36g (4.2 mmol) of [[2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester in 10 mL of methylene chloride and the mixture stirred for 1.5 hours. The mixture was concentrated at reduced pressure and residue taken up in 10 mL of methylene chloride. Anhydrous ammonia was bubbled in for 0.25 hours, the mixture stirred


-88-

for 1.5 hours and evaporated at reduced pressure. The residue was stirred with 20 mL of ethyl acetate and the mixture filtered. The filtrate was concentrated to give 1.37g of a mixture of [[3-(2-amino-1,2-dioxoethyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester and ammonium chloride. This mixture melted at 172-187°C.

**Example 2**

10 (indol-3-oxime amide starting material)


2-[[3-[[2-(Aminooxo)-1-(N-hydroxyimino)]ethyl]-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid.



A. Preparation of 2-[[3-[[2-(Aminooxo)-1-(N-hydroxyimino)]ethyl]-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetic acid methyl ester.



-89-



A stirred mixture of **1** (600 mg, 1.52 mmol) and hydroxylamine hydrochloride (528 mg, 7.60 mmol) in THF (4 mL)/CH<sub>3</sub>OH (4 mL) was heated at 55 °C for 8 h. After concentration at ambient temperature, the residue was chromatographed on silica (gradient 0-40% EtOAc in CH<sub>2</sub>Cl<sub>2</sub>) to give the title compound **2ai** (285 mg) as a white solid in 46% yield. IR (CHCl<sub>3</sub>) 3510, 3415, 1757, 1667 cm<sup>-1</sup>; <sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ 1.17 (t, *J* = 7.5 Hz, 3H), 2.84 (q, *J* = 7.5 Hz, 2H), 3.81 (s, 3H), 4.73 (s, 2H), 5.36 (s, 2H), 5.67 (br s, 1H), 6.31 (br s, 1H), 6.41 (d, *J* = 7.8 Hz, 1H), 6.87 (d, *J* = 8.2 Hz, 1H), 6.98-7.07 (m, 3H), 7.23-7.32 (m, 3H); ESIMS *m/e* 410 (*M*<sup>+</sup>+1).


Elemental Analyses for C<sub>22</sub>H<sub>23</sub>N<sub>3</sub>O<sub>5</sub>·0.30(H<sub>2</sub>O):

Calculated: C, 63.70; H, 5.73; N, 10.13;  
Found: C, 63.68; H, 5.62; N, 10.20.


### Example 3

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]glycine

-90-



**A. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]glycine methyl ester**



5


To a solution of **1** (0.100 g, 0.249 mmol) in 2 mL DMF was added collidine (0.069 mL, 0.523 mmol), methyl glycine hydrochloride (0.0313 g, 0.249 mmol), and benzotriazolyl-N-oxy-tris(dimethylamino)phosphonium hexafluorophosphate (0.115 g, 0.261) sequentially at room temperature. After 2.5 hrs. the reaction mixture was concentrated in vacuo to near dryness, then it was taken up in CH<sub>2</sub>Cl<sub>2</sub>, chromatographed on a silica gel column (gradient 20-40% THF in CH<sub>2</sub>Cl<sub>2</sub>) and dried in an 80°C vacuum oven to give 0.0768 g of **2a** as a yellow solid in 68% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t, J = 6.8 Hz, 3H), 2.90 (br q, J = 6.8 Hz, 2H),

-91-

3.57 (s, 3H), 3.88 (d,  $J = 5.5$  Hz, 2H), 4.57 (s, 2H), 5.51 (s, 2H), 6.59 (d,  $J = 5.6$  Hz, 1H), 7.01-7.08 (m 4H), 7.19-7.30 (m, 3H), 7.55 (s, 1H), 7.99 (s, 1H), 8.40 (t,  $J = 5.5$  Hz, 1H).

5

**B. Preparation of *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]glycine**




10 To a solution of 2a (0.035 g, 0.078 mmol) in 1 mL THF, 1 mL MeOH and 0.25 mL distilled H<sub>2</sub>O was added 4.17N LiOH (0.093 mL, 0.388 mmol) at room temperature. After 2 hrs. the reaction mixture was acidified with 5N HCl (0.093 mL, 0.465 mmol) and concentrated in vacuo. The residue  
15 was taken up in CH<sub>2</sub>Cl<sub>2</sub>, then rapidly triturated with hexanes to give a yellow suspension which was filtered and dried in an 80°C vacuum oven to give 0.0336 g of 3a as a yellow solid in 99% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t,  $J = 5.9$  Hz, 3H), 2.90 (br q,  $J = 5.9$  Hz, 2H), 3.80 (d,  $J = 4.8$  Hz, 2H), 4.56 (s, 2H), 5.51 (s, 2H), 6.62 (d,  $J = 5.8$  Hz, 1H),  
20 2H), 4.56 (s, 2H), 5.51 (s, 2H), 6.62 (d,  $J = 5.8$  Hz, 1H),

-92-


7.01-7.28 (m, 7H), 7.54 (s, 1H), 7.99 (s, 1H), 8.31 (t,  $J$  = 4.8 Hz, 1H), 12.25-12.75 (br s, 1H).

**Example 4**

- 5 *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-alanine



- A. Preparation of *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-alanine methyl ester
- 10



Following the experimental procedure as described for 2a, 2b was obtained as a yellow solid in 65% yield.


<sup>1</sup>H NMR (DMSO-d<sub>6</sub>)  $\delta$  1.04 (t,  $J$  = 7.2 Hz, 3H), 1.29 (d,  $J$  = 7.3 Hz, 3H), 2.91 (br q,  $J$  = 7.2 Hz, 2H), 3.54 (s, 3H), 4.29 (qd,  $J$  = 7.3, 6.8 Hz, 1H), 4.55 (s, 2H), 5.51 (s,

15

-93-

2H), 6.57 (m, 1H), 6.99 (d,  $J = 7.4$  Hz, 2H), 7.07-7.08 (m, 2H), 7.21-7.31 (m, 3H), 7.56 (s, 1H), 8.05 (s, 1H), 8.40 (d,  $J = 6.8$  Hz, 1H).

5      **B. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-alanine**




Following the experimental procedure as described for  
10 preparing compound 3a, compound 3b, was obtained as a  
yellow solid in 89% yield.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  1.04 (t,  $J$   
= 7.2 Hz, 3H), 1.29 (d,  $J = 7.3$  Hz, 3H), 2.91 (br q,  $J$   
= 7.2 Hz, 2H), 4.22 (td,  $J = 7.2, 7.1$  Hz, 1H), 4.54 (s, 2H),  
5.51 (s, 2H), 6.60 (d,  $J = 6.3$  Hz, 1H), 7.00-7.09 (m, 4H),  
15 7.21-7.30 (m, 3H), 7.53 (s, 1H), 8.03 (s, 1H), 8.31 (d,  $J$   
= 7.1 Hz, 1H), 12.75-12.84 (br s, 1H).


**Example 5**

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-  
20 indol-4-yl]oxy]acetyl]-L-leucine

-94-



A. Preparation of *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-leucine methyl ester




5

Following the experimental procedure as described for 2a, 2c was obtained as a yellow solid in 98% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.67 (d, *J* = 5.5 Hz, 3H), 0.72 (d, *J* = 5.7 Hz, 3H), 1.05 (t, *J* = 7.2 Hz, 3H), 1.30-1.50 (m, 2H), 1.51-1.64 (m, 1H), 2.91 (br q, *J* = 7.2 Hz, 2H), 3.55 (s, 3H), 4.20-4.27 (m, 1H), 4.57 (s, 2H), 5.52 (s, 2H), 6.53-6.56 (m, 1H), 6.97-7.08 (m, 4H), 7.21-7.29 (m, 3H), 7.56 (s, 1H), 8.07 (s, 1H), 8.37 (d, *J* = 7.3 Hz, 1H).

-95-

**B. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-leucine**




- 5 Following the experimental procedure as described for 3a, 3c was obtained as a yellow solid in 75% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.76 (d, J = 5.7 Hz, 3H), 0.78 (d, J = 6.1 Hz, 3H), 1.21 (t, J = 7.3 Hz, 3H), 1.39-1.43 (m, 1H), 1.69 (t, J = 7.3 Hz, 2H), 2.96 (br q, J = 7.3 Hz, 2H), 4.57-4.65 (m, 1H), 4.69 (d, J = 16.0 Hz, 1H), 4.78 (d, J = 16.0 Hz, 1H), 5.38 (s, 2H), 6.59 (d, J = 8.0 Hz, 1H), 6.89 (d, J = 8.2 Hz, 1H), 6.95-7.12 (m, 5H), 7.26-7.32 (m, 3H), 8.17 (d, J = 8.2 Hz, 1H).

15


**Example 6**

N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-aspartic acid

-96-



**A. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-aspartic acid dimethyl ester**



5

Following the experimental procedure as described for 2a, 2d was obtained as a yellow solid in 88% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t, J = 7.3 Hz, 3H), 2.72 (dd, J = 16.6, 7.1 Hz, 1H), 2.83 (dd, J = 16.7, 7.1 Hz, 1H), 2.90 (br q, J = 7.3 Hz, 2H), 3.49 (s, 3H), 3.55 (s, 3H), 4.54 (s, 2H), 4.66 (m, 1H), 5.51 (s, 2H), 6.54 (m, 1H), 6.97-7.09 (m, 4H), 7.21-7.30 (m, 3H), 7.50 (s, 1H), 7.97 (s, 1H), 8.52 (d, J = 7.9 Hz, 1H).



13-06-2001


X-12420

00944673.3(11-07-2000) - US00/16319(11-07-2000)

P DESCAMD

-97-

**B. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-aspartic acid**




5

Following the experimental procedure as described for 3a, 3d was obtained as a yellow solid in 99% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t, J = 7.2 Hz, 3H), 2.52-2.76 (m, 2H), 2.90 (br q, J = 7.2 Hz, 2H), 4.53 (s, 2H), 4.53-4.60 (m, 1H), 5.50 (s, 2H), 6.59 (d, J = 7.2 Hz, 1H), 6.98-7.09 (m, 4H), 7.19-7.30 (m, 3H), 7.47 (s, 1H), 7.94 (s, 1H), 8.33 (d, J = 7.8 Hz, 1H), 12.40-13.20 (br s, 2H).


**Example 7**

**N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-phenylalanine**

-98-



A. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-phenylalanine methyl ester




5

Following the experimental procedure as described for 2a, 2e was obtained as a yellow solid in 68% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.06 (t, J = 7.2 Hz, 3H), 2.88-3.03 (m, 4H), 3.54 (s, 3H), 4.47-4.50 (m, 1H), 4.50 (s, 2H), 5.52 (s, 2H), 6.41 (d, J = 7.7 Hz, 1H), 6.98-7.11 (m, 9H), 7.21-7.30 (m, 3H), 7.47 (s, 1H), 8.06 (s, 1H), 8.52 (d, J = 7.7 Hz, 1H).

10

-99-

**B. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-phenylalanine**




5

Following the experimental procedure as described for 3a, 3e was obtained as a yellow solid in 93% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t, J = 7.1 Hz, 3H), 2.85-3.12 (m, 4H), 4.17-4.26 (m, 1H), 4.54 (s, 2H), 5.51 (s, 2H), 6.59 (d, J = 6.4 Hz, 1H), 6.98-7.09 (m, 9H), 7.19-7.30 (m, 3H), 7.53 (s, 1H), 8.03 (s, 1H), 8.30 (d, J = 7.0 Hz, 1H), 12.50 (br s, 1H).


**Example 8**

15 [2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid

-100-



**A. Preparation of [2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid dimethyl ester**




5

Following the experimental procedure as described for 2a, 2f was obtained as a yellow solid in 98% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 1.04 (t, J = 7.3 Hz, 3H), 2.90 (br q, J = 7.3 Hz, 2H), 3.64 (s, 6H), 4.63 (s, 2H), 5.16 (d, J = 7.1 Hz, 1H), 5.51 (s, 2H), 6.54-6.56 (m, 1H), 6.98-7.09 (m, 4H), 7.21-7.30 (m, 3H), 7.43 (s, 1H), 7.88 (s, 1H), 8.90 (d, J = 7.2 Hz, 1H).

-101-

B. Preparation of [2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetamido]malonic acid




Following the experimental procedure as described for 3a,  
5 3f was obtained as a yellow solid in 99% yield. <sup>1</sup>H NMR  
(DMSO-d<sub>6</sub>) δ 1.04 (t, J = 6.9 Hz, 3H), 2.89 (br q, J = 7.3  
Hz, 2H), 4.62 (s, 2H), 4.91 (d, J = 7.2 Hz, 1H), 5.50 (s,  
2H), 6.57 (d, J = 7.2 Hz, 1H), 6.98-7.09 (m, 4H), 7.18-  
7.30 (m, 3H), 7.37 (s, 1H), 7.83 (s, 1H), 8.55 (d, J = 7.2  
10 Hz, 1H), 12.30-13.00 (br s, 2H).

-102-


Example 9

*N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1*H*-indol-4-yl]oxy]acetyl]-*L*-valine



5

A. Preparation of *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1*H*-indol-4-yl]oxy]acetyl]-*L*-valine methyl ester




10 Following the experimental procedure as described for 2a, 2g was obtained as a yellow solid in 96% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.71 (d, *J* = 6.8 Hz, 3H), 0.74 (d, *J* = 7.0 Hz, 3H), 1.05 (t, *J* = 7.2 Hz 3H), 1.99-2.05 (m, 1H), 2.90 (br q, *J* = 7.2 Hz, 2H), 3.54 (s, 3H), 4.11 (br t, *J* = 7.0 Hz, 1H), 4.60 (s, 2H), 5.52 (s, 2H), 6.52 (d, *J* = 4.4 Hz, 1H),

15

-103-

6.95 (d,  $J = 7.2$  Hz, 2H), 7.06 (br s, 2H), 7.18-7.29 (m, 3H), 7.52 (s, 1H), 8.04 (s, 1H), 8.20 (d,  $J = 7.8$  Hz, 1H).

B. Preparation of *N*-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-valine




Following the experimental procedure as described for 3a, 3g was obtained as a yellow solid in 94% yield.  $^1\text{H}$  NMR (DMSO- $d_6$ )  $\delta$  0.71 (d,  $J = 6.9$  Hz, 3H), 0.75 (d,  $J = 6.8$  Hz, 3H), 1.04 (t,  $J = 7.3$  Hz, 3H), 2.01-2.07 (m, 1H), 2.90 (br q,  $J = 7.3$  Hz, 2H), 4.09 (br dd,  $J = 7.9, 6.2$  Hz, 1H), 4.60 (s, 2H), 5.51 (s, 2H), 6.54 (d,  $J = 6.1$  Hz, 1H), 6.95 (d,  $J = 7.3$  Hz, 2H), 6.99-7.08 (m, 2H), 7.18-7.29 (m, 3H), 7.49 (s, 1H), 8.01 (s, 1H), 8.08 (d,  $J = 8.2$  Hz, 1H), 12.63 (br s, 1H).

-104-


**Example 10**

**N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine**



5

**A. Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine methyl ester**




10 Following the experimental procedure as described for 2a, 2h was obtained as a yellow solid in 73% yield. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ 0.64-0.71 (m, 6H), 0.99-1.08 (m, 4H), 1.21-1.26 (m, 1H), 1.76-1.80 (m, 1H), 2.91 (br q, J = 7.4 Hz, 2H), 3.53 (s, 3H), 4.15 (br t, J = 7.2 Hz, 1H), 4.60 (s,



-105-

2H), 5.52 (s, 2H), 6.52 (m, 1H), 6.96 (d,  $J = 7.2$  Hz, 2H),  
7.02-7.07 (m, 2H), 7.18-7.29 (m, 3H), 7.53 (s, 1H), 8.04  
(s, 1H), 8.23 (d,  $J = 7.7$  Hz, 1H).

5            B.    Preparation of N-[2-[[3-(Aminooxoacetyl)-2-ethyl-1-(phenylmethyl)-1H-indol-4-yl]oxy]acetyl]-L-isoleucine



Following the experimental procedure as described for 3a,  
10 3h was obtained as a yellow solid in 92% yield.  $^1\text{H}$  NMR  
(DMSO- $d_6$ )  $\delta$  0.64-0.84 (m, 6H), 1.04 (t,  $J = 7.2$  Hz, 3H),  
1.21-1.28 (m, 2H), 1.76-1.80 (m, 1H), 2.91 (br q,  $J = 7.2$   
Hz, 2H), 4.12 (br t,  $J = 7.3$  Hz, 1H), 4.59 (s, 2H), 5.51  
(s, 2H), 6.55 (d,  $J = 6.4$  Hz, 1H), 6.96 (d,  $J = 7.2$  Hz,  
15 2H), 7.01-7.08 (m, 2H), 7.21-7.29 (m, 3H), 7.51 (s, 1H),  
8.01 (s, 1H), 8.11 (d,  $J = 7.4$  Hz, 1H), 12.40-12.65 (br s,  
1H).

-106-

**Assay**

The following chromogenic assay procedure was used to identify and evaluate inhibitors of recombinant human secreted phospholipase A<sub>2</sub>. The assay described herein

5 has been adapted for high volume screening using 96 well microtiter plates. A general description of this assay method is found in the article, "Analysis of Human Synovial Fluid Phospholipase A<sub>2</sub> on Short Chain

10 Phosphatidylcholine-Mixed Micelles: Development of a Spectrophotometric Assay Suitable for a Microtiterplate Reader", by Laure J. Reynolds, Lori L. Hughes, and Edward A Dennis, Analytical Biochemistry, 204, pp. 190-197, 1992 (the disclosure of which is incorporated herein by reference):

## 15 Reagents:

## REACTION BUFFER -

CaCl<sub>2</sub>·2H<sub>2</sub>O (1.47 g/L)

KCl (7.455 g/L)

Bovine Serum Albumin (fatty acid free) (1 g/L)

20 (Sigma A-7030, product of Sigma Chemical Co., St. Louis MO, USA)

TRIS HCl (3.94 g/L)

pH 7.5 (adjust with NaOH)

## ENZYME BUFFER -

25 0.05 NaOAc.3H<sub>2</sub>O, pH 4.5

0.2 NaCl

-107-

Adjust pH to 4.5 with acetic acid

DTNB - 5,5'-dithiobis-2-nitrobenzoic acid

RACEMIC DIHEPTANOYL THIO - PC

5 racemic 1,2-bis(heptanoylthio)-1,2-dideoxy-*sn*-  
glycero-3-phosphorylcholine

TRITON X-100<sup>TM</sup> prepare at 6.249 mg/ml in  
reaction buffer to equal 10uM.

REACTION MIXTURE -

10 A measured volume of racemic dipheptanoyl thio PC  
supplied in chloroform at a concentration of 100  
mg/ml is taken to dryness and redissolved in 10  
millimolar

15 TRITON X-100<sup>TM</sup> nonionic detergent aqueous solution.  
Reaction Buffer is added to the solution, then DTNB  
to give the Reaction Mixture.

The reaction mixture thus obtained contains 1mM  
diheptanoly thio-PC substrate, 0.29 mM Triton X-  
100<sup>TM</sup> detergent, and 0.12 mM DTMB in a buffered  
aqueous solution at pH 7.5.

20

Assay Procedure:

1. Add 0.2 ml reaction mixture to all wells;
2. Add 10 ul test compound (or solvent blank) to  
appropriate wells, mix 20 seconds;
- 25 3. Add 50 nanograms of sPLA<sub>2</sub> (10 microliters) to  
appropriate wells;

-108-

4. Incubate plate at 40 °C for 30 minutes;
5. Read absorbance of wells at 405 nanometers with an automatic plate reader.

5 All compounds were tested in triplicate. Typically, compounds were tested at a final concentration of 5 ug/ml. Compounds were considered active when they exhibited 40% inhibition or greater compared to uninhibited control reactions when measured

10 at 405 nanometers. Lack of color development at 405 nanometers evidenced inhibition. Compounds initially found to be active were reassayed to confirm their activity and, if sufficiently active, IC<sub>50</sub> values were determined. Typically, the IC<sub>50</sub> values (see, Table I,

15 below) were determined by diluting test compound serially two-fold such that the final concentration in the reaction ranged from 45 ug/mL to 0.35 ug/ml. More potent inhibitors required significantly greater dilution. In all cases, % inhibition measured at 405

20 nanometers generated by enzyme reactions containing inhibitors relative to the uninhibited control reactions was determined. Each sample was titrated in triplicate and result values were averaged for plotting and calculation of IC<sub>50</sub> values. IC<sub>50</sub> were determined by

25 plotting log concentration versus inhibition values in the range from 10-90% inhibition.

-109-

Results of Human Secreted Phospholipase A<sub>2</sub> Inhibition  
Tests

Table

Compound No. from Examples 3-10	Inhibition of human secreted PLA <sub>2</sub> IC <sub>50</sub> ± mean deviation (3-4 tests) (nM)
1	49
2A	529
2B	533
2C	82
2D	874
2E	666
2F	698
2G	283
2H	166
3A	71
3B	59
3C	28
3D	132
3E	64
3F	44.7
3G	36.4
3H	25.1

5        The compound of Example 1 is highly active in  
inhibiting sPLA<sub>2</sub>.

While the present invention has been illustrated  
above by certain specific embodiments, it is not intended  
10    that these specific examples should limit the scope of the  
invention as described in the appended claims.